
# in September **Routt County** Common Birds of



& other local photographers Photos by Yampa Valley Birding Club

Note: Descriptions are of breeding males (R): Resident; (M): Migratory

[ www.coloradocranes.org ]



PERCHING BIRDS

# ☐ Brewer's Blackbird (M) glossy black overall, long tail, yellow eye fields, open areas

red wing patch marshes, wet fields black overall with Blackbird (R) □ Red-winged



open areas with trees pale blue overall

brushy understory of woods ☐ **Gray Catbird (M)** gray overall with black cap

rufous undertail coverts



black cap and throat, white cheeks, buffy flanks □ Black-capped Chickadee (R)



dark grey overall, short tail ☐ American Dipper (R) mountain streams

suburban habitats pale overall with dark collar

Dove (R)



☐ Mourning Dove (M) gray-brown overall, long, pointed tail open areas

orange-red cap, throat

☐ House Finch (R)

breast; brownish back brushy, wooded areas

# PERCHING BIRDS

bright yellow overall, black ☐ American Goldfinch (R) orchards, overgrown fields wings and forehead



☐ Evening Grosbeak (R) large bill, yellow and mixed forests black body



☐ **Stellar's Jay (R)** dark blue body, black coniferous and mixed head, long crest

☐ American Crow (R)
black overall

open habitats



☐ Dark-eyed Junco (R) gray overall, black lores, open woods rufous back

blue-gray above, white below; dark breast band, ☐ Belted Kingfisher (R) sheltered open water shaggy crest



☐ Lesser Goldfinch (M) yellow below, greenish above with black cap patchy open habitats



rosy red throat, green body dry coniferous woods Hummingbird (M) □ Broad-tailed



wooded areas

below; gray cheek and back ☐ Western Scrub Jay (R) dark blue above, whitish





bold face pattern, dark mask mostly barren open areas brown above, dark breast band ☐ Horned Lark (R)



streaked brown with yellow on wings, chest, tail ☐ Pine Siskin (R) various habitats

☐ House Sparrow (R) open forests

black throat and breast; gray head and belly rufous-brown above, cities, towns



PERCHING BIRDS

☐ Black-billed Magpie (R) bold black and white white wing patches long dark tail,



Nuthatch (R) blue above, white below, white face, black crown ☐ White-breasted gray overall; dark head neck, breast; black wing bars, ☐ Rock Pigeon (R) widespread, various



☐ **Common Raven (R)**black; large bill,
wedge-shaped tail ☐ American Robin (R) gray above, rusty below open woodlands



brown above, gray below ☐ Chipping Sparrow (M)

□ Violet-green



eyebrow, dark eye-line open woodlands □Wilson's Warbler (M) woodlands, urban areas yellow overall

gray eyebrow, brown crown streaked yellow breast ☐ Lincoln's Sparrow (M) dense brushy areas

olive above, black cap

brushy woods

open forests and edges



brown above, yellow below, black breast band brown above, streaky breast ☐ Savannah Sparrow (R) open grasslands, marshes dark mustache

brushy areas near water

☐ **Song Sparrow (R)** brown streaks on head

rufous-tinged wings,

streaked breast

Meadowlark (R)

□ Western

grasslands



☐ Barn Swallow (M) dark blue above, cinnamon

open areas, esp. near water

below; rusty brown forehead and throat



chestnut throat, white forehead dark blue above, white below ☐ Cliff Swallow (M) open riparian areas ☐ Tree Swallow (M) blue-green above,

dark metallic





bluish overall; bright yellow throat, flanks, rump ☐ Yellow-rumped Warbler (M)

mask; yellow-tipped tail ☐ Cedar Waxwing (R) silky brown with black brown crest bushy areas



Wood-pewee (M) gray overall, dark head 2 dull wing bars, mature forests and edges



WOODPECKERS

# wooded areas with openings ☐ Red-shafted Flicker (R) Black breast band, barred back, spotted belly



white back, white below black wings, white and black stripes on head wooded habitats

dense brushy habitats

gray-brown overall ☐ House Wren (M)

barring above; pale eyebrow



☐ Downy Woodpecker (R) ☐ Hairy Woodpecker (R) Similar to Downy but much larger with larger bill

mature forests



□Great Blue Heron (M) black crown stripe on whitish head large, gray overall, open habitats



below, spots on chest, teeters body constantly ☐ Spotted Sandpiper (M) brown above, white



brown above, dark bars c ☐ Greater Yellowlegs (1 flanks, long yellow legs, long bill mudflats, shallow water

brownish overall, cream-colored stripes on back striped head, long bill


[www.coloradocranes.org]

☐ Wilson's Snipe (R)

marshes, damp fields



deep lakes and rivers





gray overall, often rust-stained, red crown, white face open meadows, marshes, fields ☐ Sandhill Crane (M)



band, rufous rump and tail below, dark double breast brown above, white ☐ Killdeer (M) open ground





□ Common Merganser (M) □ Sharp-shinned Hawk (R) With dark breast; white white; black back, dark green head, orange bill reddish below; small with constructions of the sharp shinned Hawk (R) with dark breast; white white white white white with face; white underside with the sharp shinned Hawk (R) with dark breast; white wh





brown above, pale breast white cheek, dark bill open ground near water ☐ Canada Goose (R) black neck,



blue-gray above, pale ☐ Cooper's Hawk (M) woods and edges long banded tail reddish below,



☐ Swainson's Hawk (M)

dark above; pale belly with dark breast; white open country



BIRDS OF PREY

BIRDS OF PRES

brown body and wings white head and tail, ☐ Bald Eagle (R)

☐ American Kestral (R)

rufous above,

blue-gray wings, 2 dark streaks on head

open country

dark above, white below,

Osprey (M)

white cap with dark



white rump, black wingtips gray above, whitish below, ☐ Northern Harrier (R) open country, wetlands



red tail, brown above, whitish belly with dark band ☐ Red-Tailed Hawk (R) various habitats; edges ☐ Great Horned Owl (R) grayish above, barred



nocturnal, various habitats below, large ear tufts







